

This is a repository copy of *Measurement of the 92,93,94,100Mo(γ ,n) reactions by Coulomb Dissociation*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/102561/>

Version: Published Version

Article:

Göbel, K., Adrich, P., Altstadt, S. et al. (70 more authors) (2016) Measurement of the 92,93,94,100Mo(γ ,n) reactions by Coulomb Dissociation. Journal of Physics: Conference Series. 012034. ISSN 1742-6596

<https://doi.org/10.1088/1742-6596/665/1/012034>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Measurement of the $^{92,93,94,100}\text{Mo}(\gamma, n)$ reactions by Coulomb Dissociation

This content has been downloaded from IOPscience. Please scroll down to see the full text.

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 144.32.224.203

This content was downloaded on 15/07/2016 at 17:27

Please note that [terms and conditions apply](#).

Measurement of the $^{92,93,94,100}\text{Mo}(\gamma, n)$ reactions by Coulomb Dissociation

K Göbel¹, P Adrich², S Altstadt¹, H Alvarez-Pol³, F Aksouh², T Aumann², M Babilon⁴, K-H Behr², J Benlliure³, T Berg⁵, M Böhmer⁶, K Boretzky², A Brünle², R Beyer⁷, E Casarejos⁸, M Chartier⁹, D Cortina-Gil², A Chatillon³, U Datta Pramanik¹⁰, L Deveau¹¹, M Elvers^{4,12}, T W Elze¹, H Emling², M Erhard⁷, O Ershova^{1,2}, B Fernandez-Dominguez⁹, H Geissel², M Górski², T Heftrich¹, M Heil², M Hellstroem², G Ickert², H Johansson^{2,13}, A R Junghans⁷, F Käppeler¹⁴, O Kiselev⁵, A Klimkiewicz², J V Kratz⁵, R Kulesa¹⁵, N Kurz², M Labiche¹⁶, C Langer^{1,2}, T Le Bleis^{2,17}, R Lemmon¹⁸, K Lindenberg⁴, Y A Litvinov², P Maierbeck⁶, A Movsesyan^{2,4}, S Müller⁴, T Nilsson¹³, C Nociforo², N Paar¹⁹, R Palit², S Paschalis⁹, R Plag^{1,2}, W Prokopowicz², R Reifarth^{1,2}, D M Rossi^{2,5}, L Schnorrenberger⁴, H Simon², K Sonnabend¹, K Sümmerer², G Surówka¹⁵, D Vretenar¹⁹, A Wagner⁷, S Walter¹⁴, W Waluś¹⁵, F Wamers², H Weick², M Weigand¹, N Winckler¹⁴, M Winkler² and A Zilges^{4,12}

¹ Goethe-Universität Frankfurt a. M., Germany

² GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany

³ Universidad de Santiago de Compostela, Spain

⁴ Technische Universität Darmstadt, Germany

⁵ Johannes Gutenberg-Universität Mainz, Germany

⁶ Technische Universität München, Germany

⁷ Helmholtz-Zentrum Dresden-Rossendorf, Germany

⁸ Universidade de Vigo, Spain

⁹ University of Liverpool, United Kingdom

¹⁰ SINP Kolkata, India

¹¹ Université de Paris Sud, Orsay, France

¹² Institut für Kernphysik, Universität zu Köln, Germany

¹³ Chalmers University of Technology, Göteborg, Sweden

¹⁴ FZ Karlsruhe, Germany

¹⁵ Jagellonian University Krakow, Poland

¹⁶ University of Paisley, United Kingdom

¹⁷ University of Strasbourg, France

¹⁸ CCLRC Daresbury Laboratory, United Kingdom

¹⁹ University of Zagreb, Croatia

E-mail: goebel@physik.uni-frankfurt.de



Abstract. The Coulomb Dissociation (CD) cross sections of the stable isotopes $^{92,94,100}\text{Mo}$ and of the unstable isotope ^{93}Mo were measured at the LAND/R³B setup at GSI Helmholtzzentrum für Schwerionenforschung in Darmstadt, Germany. Experimental data on these isotopes may help to explain the problem of the underproduction of $^{92,94}\text{Mo}$ and $^{96,98}\text{Ru}$ in the models of p-process nucleosynthesis. The CD cross sections obtained for the stable Mo isotopes are in good agreement with experiments performed with real photons, thus validating the method of Coulomb Dissociation. The result for the reaction $^{93}\text{Mo}(\gamma, n)$ is especially important since the corresponding cross section has not been measured before. A preliminary integral Coulomb Dissociation cross section of the $^{94}\text{Mo}(\gamma, n)$ reaction is presented. Further analysis will complete the experimental database for the (γ, n) production chain of the p-isotopes of molybdenum.

1. Motivation

The p-nuclei between ^{74}Se and ^{196}Hg are produced under explosive conditions in a sequence of photodissociations of s- and r-process seeds and subsequent β -decays [1]. The modeling of the p-process requires a large network with more than 2,000 isotopes linked by more than 20,000 reactions. The necessary nuclear physics input includes masses of the isotopes, half-lives of the nuclei involved, and reaction rates. Only a small fraction of the required reaction rates can be determined experimentally, most of which rely on predictions from statistical model calculations [2]. An experimental validation of the reaction rates included in stellar models is therefore highly desired.

$^{92,94}\text{Mo}$ and $^{96,98}\text{Ru}$ are the most abundant, but not sufficiently explained p-nuclei [2]. According to recent stellar model calculations, ^{94}Mo is mainly synthesized via the (γ, n) photodisintegration chain starting from the more neutron-rich, and stable molybdenum isotopes [3] (Fig. 1). In order to understand the abundance ratio of ^{94}Mo to ^{92}Mo , the determination of the cross section of the $^{94}\text{Mo}(\gamma, n)$ reaction is necessary. If this reaction is possible in a certain stellar environment, the photodisintegration of ^{93}Mo will follow immediately due to the lower neutron separation energy (Fig. 2). The closed neutron shell at ^{92}Mo terminates the photoneutron chain of the molybdenum isotopes.




Figure 1. Portion of the reaction network leading to the production of the p-isotopes ^{92}Mo and ^{94}Mo .




Figure 2. Neutron separation energies of the molybdenum isotopes ^{92}Mo , ^{93}Mo , and ^{94}Mo .

The $^{92,93,94,100}\text{Mo}(\gamma, n)$ reactions were measured via Coulomb Dissociation at the LAND/R³B setup at GSI Helmholtzzentrum für Schwerionenforschung in Darmstadt, Germany. The (γ, n) reactions are studied in inverse kinematics. The nuclei under investigation are prepared as an ion beam, allowing radioactive nuclei to be investigated. The aim of the present experiment was the validation of the method of Coulomb Dissociation by comparing the results with data from

photoactivation measurements [4]. The (γ, n) cross section of the unstable isotope ^{93}Mo could be determined for the first time [5].

2. Experimental method

Most nuclei involved in photodissociation reactions in stellar nucleosynthesis networks are unstable and cannot be prepared as a sample for experiments using real photons. One solution is to study the (γ, n) reaction in inverse kinematics: The nucleus under investigation hits a high-Z target and interacts with the time-varying Coulomb field. This interaction can be interpreted as an absorption of a virtual photon (Fig. 3) [6]. The absorption probability can be translated into a (γ, n) cross section.




Figure 3. Schematic view of the Coulomb excitation process of a molybdenum nucleus. The Mo projectile impinges on a lead target. The target's electromagnetic field \vec{E} seen by the projectile is Lorentz-contracted in the direction of motion. In a peripheral collision (large impact parameter b) the projectile can be excited by a virtual photon.

The experimental setup is shown in Fig. 4. The ion beam enters the experimental area and is tracked in time by the POS scintillation detector. Position sensitive pin diodes (PSP) [7] detect the position of the ion as well as its charge via energy loss in the detector. The heavy fragment is deflected by the ALADiN magnet (A Large Acceptance Dipole magNet) after the (γ, n) reaction and hits three scintillating fibre detectors (GFI) [8] where the horizontal position is measured. The TFW (Time-of-Flight Wall) at the end of the fragment arm provides position, time and charge information. The emitted neutron remains unaffected by the magnetic field and is detected by the Large Area Neutron Detector (LAND) [9].

3. Analysis and preliminary results

Several steps are needed to extract the cross section from the data of the complex experimental setup.

3.1. Selection of the reaction channel

The reaction channel $^{94}\text{Mo}(\gamma, n)^{93}\text{Mo}$ has to be selected by several conditions applied to the data. A primary beam of ^{94}Mo entered the experimental area. To reject breakup events in the material in front of the target, the incoming projectile ^{94}Mo is identified by the charge (Z) measured by PSP1 and PSP2. In the outgoing channel, the heavy reaction fragment must be identified by its charge and mass. The fragment charge is selected by data obtained by the PSP3 and the TFW. The fragment mass is determined by tracking of the ions through the magnetic field, using position information provided by the PSP and GFI detectors. Furthermore, one neutron has to be detected by LAND.




Figure 4. LAND/R³B experimental setup for the measurement of $^{92,93,94,100}\text{Mo}(\gamma, n)$ by Coulomb Dissociation [10].

3.2. Background subtraction

In addition to Coulomb excitation, the projectile may experience nuclear interactions in the target and reactions with the material outside of the target. In order to determine these background contributions, two additional measurements were performed: one without target (“empty target”) and one with a carbon target. Due to the lower atomic number of carbon, the induced electromagnetic excitations are much weaker compared to lead. The cross section of the nuclear interaction is related to the size (radii) of the projectile and of the target nuclei. Therefore, the interaction of the projectile with the carbon target is considered to be purely nuclear [11]. The background distributions are normalized and subtracted from the lead target distributions. The Coulomb Dissociation cross section σ_{CD} is determined by

$$\sigma_{CD} = \left(\frac{M_{Pb}}{d_{Pb}N_A} \right) p_{Pb} - \left(\alpha \cdot \frac{M_C}{d_CN_A} \right) p_C - \left(\frac{M_{Pb}}{d_{Pb}N_A} - \alpha \cdot \frac{M_C}{d_CN_A} \right) p_{Em}, \quad (1)$$

where p is the interaction probability, M the molar mass of the target material [g mol^{-1}], d the target thickness, N_A the Avogadro number [mol^{-1}] and α the nuclear scaling factor. The indices refer to the different targets. In the analysis presented here, α is calculated according to the “black disc model” [12], in which the interacting nuclei are considered fully opaque to each other.

$$\alpha = \frac{A_P^{\frac{1}{3}} + A_{Pb}^{\frac{1}{3}}}{A_P^{\frac{1}{3}} + A_C^{\frac{1}{3}}}. \quad (2)$$

Here, A_P , A_{Pb} , and A_C represent the mass numbers of the projectile, the lead and the carbon target, respectively. The model yields $\alpha = 1.53$ for the ^{94}Mo projectile.

3.3. Preliminary integrated Coulomb Dissociation cross section

For the determination of the integrated Coulomb Dissociation cross section, a distribution of any observable derived according to Eq. 1 can be used. In the preliminary analysis presented

here, the neutron kinetic energy in the fragment rest frame E_n was chosen. It is calculated on an event-by-event basis with the kinematics of the incoming ion ^{94}Mo , the outgoing fragment ^{93}Mo and the evaporated neutron considering the Q value of the reaction:

$$E_n = \sqrt{m_{93\text{Mo}}^2 + m_n^2 + 2 \cdot \gamma_{93\text{Mo}} \gamma_n \cdot m_{93\text{Mo}} m_n \cdot (1 - \beta_{93\text{Mo}} \beta_n \cos \theta_{93\text{Mo},n}) c^2 - m_{94\text{Mo}} c^2 - Q}. \quad (3)$$

The spectrum was corrected for the LAND efficiency and acceptance. The LAND efficiency and acceptance as a function of the neutron kinetic energy was simulated considering the nominal efficiency of LAND, which was determined in a calibration experiment, the inefficiency resulting from switched-off paddles, and the limited geometrical acceptance of the detector. The data were taken from Ref. [10].

The derived spectrum is shown in Fig. 5. Its integral delivers a preliminary integrated Coulomb Dissociation cross section of 675 mb for the reaction $\text{Pb}(^{94}\text{Mo}, ^{93}\text{Mo}+n)\text{Pb}$ at a beam energy of 500 MeV.




Figure 5. Preliminary Coulomb Dissociation cross section of the reaction $\text{Pb}(^{94}\text{Mo}, ^{93}\text{Mo}+n)\text{Pb}$ at a beam energy of 500 MeV. The errors indicated are purely statistical.

4. Comparison to results from photoabsorption measurements

The results for the reactions $^{92}\text{Mo}(\gamma, n)$ and $^{100}\text{Mo}(\gamma, n)$ are in good agreement with measurements performed with real photons [5]. According to the method described in [10], the photoabsorption data available for the reaction $^{94}\text{Mo}(\gamma, n)^{93}\text{Mo}$ from Beil *et al.* [13] were converted to Coulomb excitation spectra for the E1 and E2 components using the systematics for Giant Quadrupole Resonances. The data are presented in Fig. 6.

From the photoabsorption data, a Coulomb Dissociation cross section of 909 mb is calculated. The ratio of the integral and of the integrated cross section deduced from the experimental data was found to be 0.74. As a first approximation, the obtained value agrees with the normalization factor suggested by Berman *et al.* [14] and Erhard *et al.* [15], that needs to be applied to the photoabsorption data by Beil *et al.* The factor was confirmed in Ref. [10] for ^{92}Mo and ^{100}Mo . The results of further analysis of the data for the $^{94}\text{Mo}(\gamma, n)^{93}\text{Mo}$ reaction, of additional efficiency corrections and of a detailed study of the systematical errors will provide a full picture of the Coulomb Dissociation experiment of the $^{92,93,94,100}\text{Mo}(\gamma, n)$ reactions at the LAND/R³B setup.

5. Summary and outlook

The Coulomb Dissociation (CD) cross sections of the stable isotopes $^{92,94,100}\text{Mo}$ and of the unstable isotope ^{93}Mo were measured at the LAND/R³B setup at GSI Helmholtzzentrum für Schwerionenforschung in Darmstadt, Germany. The CD cross sections obtained for ^{92}Mo and




Figure 6. Coulomb excitation cross section of ^{94}Mo derived from the data of Beil *et al.* [13]. The data were obtained by a convolution of the E1 and E2 photoabsorption cross sections with the corresponding calculated virtual photon field. The E2 component was determined from the systematics for Giant Quadrupole Resonances [10].

^{100}Mo and the preliminary value for ^{94}Mo are in good agreement with experiments using real photons, thus validating the method of Coulomb Dissociation. The result of $^{93}\text{Mo}(\gamma, n)$ is particularly important since the corresponding cross section has not been measured before. The results from the ongoing analysis of $^{94}\text{Mo}(\gamma, n)^{93}\text{Mo}$ will complete the analysis of this series of measurements, hence completing the experimental database for the (γ, n) production chain of the p-isotopes of molybdenum.

Acknowledgments

This project was supported by the Helmholtz International Center for FAIR, the Helmholtz Young Investigator Group VH-NG-327, DFG (SO907/2-1) and HGS-HiRe.

References

- [1] Lambert D L 1992 *The Astronomy and Astrophysics Review* **3** 201–256
- [2] Arnould M and Goriely S 2003 *Physics Reports* **384** 1–84
- [3] Travaglio C, Röpke F K, Gallino R and Hillebrandt W 2011 *The Astrophysical Journal* **739** 93
- [4] Sonnabend K, Babilon M, Hasper J, Müller S, Zarza M and Zilges A 2006 *European Physical Journal A* **27** 149–152
- [5] Ershova O *et al* 2010 Coulomb dissociation reactions on Mo isotopes for astrophysics applications *Nuclei in the Cosmos*
- [6] Bertulani C A and Baur G 1988 *Physics Reports* **163** 299–408
- [7] Mahata K *et al* 2009 *Nuclear Instruments and Methods in Physics Research A* **608** 331–335
- [8] Cub J *et al* 1998 *Nuclear Instruments and Methods in Physics Research A* **402** 67–74
- [9] Blaich *et al* 1992 *Nuclear Instruments and Methods in Physics Research A* **314** 136–154
- [10] Ershova O 2012 *Coulomb Dissociation Reactions on Molybdenum Isotopes for Astrophysics Applications* Ph.D. thesis Goethe University Frankfurt, Germany
- [11] Bertulani C A and Gade A 2010 *Physics Reports* **485** 195–259
- [12] Aumann T, Kratz J V, Stiel E, Sümmerer K, Brüche W, Schädel M, Wirth G, Fauerbach M and Hill J C 1993 *Physical Review C* **47** 1728–1737
- [13] Beil H, Bergère R, Carlos P, Leprêtre A, Miniac A D and Veyssière A 1974 *Nuclear Physics A* **227** 3
- [14] Berman B L and Fultz S C 1975 *Reviews of Modern Physics* **47** 3
- [15] Erhard M, Junghans A R, Nair C, Schwengner R, Beyer R, Klug J, Kosev K, Wagner A and Grosse E 2010 *Physical Review C* **81** 034319